

**BEAMLINE**  
X18B

**PUBLICATION**

R.K. Egdal, A. Hazell, F.B. Larsen, C.J. McKenzie, and R.C. Scarrow, "A Dihydroxo-bridged Fe(II)-Fe(III) Complex: A New Member of the Diiron Diamond Core Family." *J. Am. Chem. Soc.*, **125**, 32-33 (2003).

**FUNDING**

DANSYNC  
Seimens Fund (CJM, FBL)

**FOR MORE INFORMATION**

Christine McKenzie, University of Southern Denmark  
chk\_chem@chem.sdu.dk

## Electron Localization in a Mixed-Valent (+2/+3) Di-Iron Complex


Frank B. Larsen<sup>1</sup>, Christine J. McKenzie<sup>1</sup>, and Robert C. Scarrow<sup>2</sup>

<sup>1</sup> University of Southern Denmark, Odense Campus; <sup>2</sup>Haverford College

*Mixed-valence coordination complexes are molecules containing at least two of the same metal ions, where the formal oxidation state (charge) of the metal ions is different. Such complexes are of interest to inorganic chemists because of their unusual electronic and magnetic properties. An average (sometimes non-integral) charge may be the best assignment for oxidation state due to delocalization of electrons, such that the charge at each metal ion is effectively the same value. Accurate determination of bond lengths can be used to assess the extent of electronic delocalization. X-ray absorption spectroscopy (XAS) performed at the NSLS was used to determine that a mixed-valence complex with a  $Fe_2(OH)_2$  core is best described as a complex with one  $Fe^{2+}$  and one  $Fe^{3+}$  ion. The XAS results were used along with X-ray diffraction to determine the crystal structure.*


Dinuclear iron complexes containing a  $Fe_2O_2$  rhombic core are of interest as models of the structure and function of the active site of the class of non-heme iron enzymes to which the hydroxylation protein of methane monooxygenase (MMOH) belongs. This enzyme is responsible for bacterial conversion of methane to methanol. Recently, a mixed valence ( $Fe^{II}Fe^{III}$ ) form of MMOH was structurally characterized as containing a triply-protonated rhombic core (i.e. one  $\mu$ -OH and one  $\mu$ -OH<sub>2</sub> bridging between the iron atoms). [Wittington, D. A.; Lippard, S. J. *J. Am. Chem. Soc.* 2001, **123**, 827-838.] While synthetic  $Fe_2O_2(H)_{0-2}$  compounds with a rhombic core exist for the II-II, III-III, III-IV oxidation states, the II-III state was unknown.

Using the neutral capping ligand, *N,N'*-dimethyl-*N,N'*-bis(2-pyridylmethyl)ethane-1,2-diamine (L), to complete the coordination requirements of the iron atoms, and thus prevent an uncontrolled polymerization to form iron(III) hydroxides, McKenzie and coworkers at the University of Southern Denmark prepared and crystallized the first example of a synthetic diiron complex in the II-III iron oxidation state and containing a



A  $Fe_2O_2(H)_{0-4}$  rhombic core.

rhombic core. The compound,  $[L_2Fe_2(OH)_2](ClO_4)_3 \cdot 3H_2O$ , is unstable in solution. In the solid state, crystals decompose in weeks. The products of decomposition are iron (III) compounds. The initial structure of  $[L_2Fe_2(OH)_2](ClO_4)_3 \cdot 3H_2O$ , determined by single crystal X-ray diffraction, showed that the two iron atoms were related by sym-



Neutral capping ligand, *N,N'*-dimethyl-*N,N'*-bis(2-pyridylmethyl)ethane-1,2-diamine

metry, suggesting a delocalized mixed-valence complex ( $Fe^{+2.5}Fe^{+2.5}$ ) with an average Fe-O distance of 1.98Å. All other things being equal, the metal-donor distances depend on metal oxidation state and are shorter for higher oxidation states. Given the chemical similarity of the local environment of each iron atom of the molecule, identical oxidation states (of +2.5) seemed plausible. The physical interpretation of such a non-integer oxidation state is that there is little barrier for electron transfer between the metal ions. However an alternative explanation is that the compound can be regarded as a valence trapped  $Fe^{II}$ - $Fe^{III}$  compound and the crystal



Frank B. Larsen and Christine J. McKenzie at NSLS Beamline X18B.


structure is disordered.

In experiments performed at Beamline X18B at the NSLS, Larsen, McKenzie, and Scarrow found that structural disorder in the crystal lattice is indeed the cause of the apparent equivalent oxidation states. The best fit of the X-ray absorption fine structure (XAFS) measurements was obtained by refining 2 Fe-O distances each to  $1.90 \pm 0.03$  and  $2.12 \pm 0.08$  Å rather than using the Fe-O distance found in the crystal structure (**Figure 1**). The refined Debye-Waller factor ( $\sigma^2$ ) for these Fe-O shells was  $0.004$  Å<sup>2</sup>, typical of what is found for compounds with only vibrational disorder in the bond


lengths. The iron K-edge (XANES) spectrum could be well modelled as arising from 50% Fe<sup>2+</sup> and 50% Fe<sup>3+</sup> (**Figure 2**). The combined results from XANES and EXAFS analysis indicate that the iron atoms in  $[L_2Fe_2(OH)_2](ClO_4)_3 \cdot 3H_2O$  are localized in their valence state on the X-ray time scale. Since the same time scale applies to X-ray diffraction, the apparent equivalence in Fe-O bond lengths in the crystal structure is ascribed to packing disorder.

Based on these results, the single crystal diffraction data of  $[L_2Fe_2(OH)_2](ClO_4)_3 \cdot 3H_2O$  were re-fit using two half-occupancy isotropic oxygen atoms instead of the

single anisotropic bridging oxygen atom (**Figure 3**). The two models fit the diffraction data equally well ( $R_w = 0.044$ ) but the disordered model is obviously preferred because it explains the EXAFS and XANES results. In the disordered model, the averaged distances ascribed to Fe<sup>III</sup>-O ( $1.89$  Å) and to Fe<sup>II</sup>-O ( $2.07$  Å), as well as the Fe-Fe distance ( $2.95$  Å), are within the uncertainty ranges for these parameters established by EXAFS analysis. These results show that, despite an ostensibly symmetrical chemical environment for both iron atoms of  $[L_2Fe_2(OH)_2](ClO_4)_3 \cdot 3H_2O$ , the electrons localize such that one Fe<sup>II</sup> and one Fe<sup>III</sup> are present in the complex.



**Figure 1.** EXAFS analysis of  $[L_2Fe_2(OH)_2](ClO_4)_3 \cdot 3H_2O$ . (A) The fluorescence (circles) and transmission (diamond) detected EXAFS are compared (both are offset by  $+4$  Å<sup>-3</sup>; no systematic difference are evident). (B) Fourier filtered EXAFS (thick line) and simulation of the EXAFS to determine Fe-X distances discussed in this article.



**Figure 2.** XANES of  $[L_2Fe_2(OH)_2](ClO_4)_3 \cdot 3H_2O$  and its simulation representing contributions of 50% from Fe<sup>II</sup> and 50% from Fe<sup>III</sup> XANES shapes.

**Figure 3.** X-ray crystal structure of the cation of  $[L_2Fe_2(OH)_2](ClO_4)_3 \cdot 3H_2O$ , with enlarged geometry of the  $Fe_2O_2H_2$  rhombic core inferred from a combination of XAS and X-ray diffraction experiments.

